



\*Continued from Directions for Use

#### FOR CANINE PARVOVIRUS ACTIVITY:

KennelSol HC is effective against Canine Parvovirus at 2 oz. per gallon. Follow Veterinary Practice / Animal Care / Animal Laboratory / Zoos / Pet Shop / Kennels Disinfection Directions.

#### FARM PREMISE, LIVESTOCK, POULTRY & TURKEY HOUSE DISINFECTANT

DILUTION: 1:256 1/2 OUNCE PER GALLON OF WATER

1. Remove all animals and feeds from premises, trucks, coops, crates, and enclosures.
2. Remove all litter and manure from floors, walls, and surfaces of barns, pens, stalls, chutes, vehicles, and other facilities and fixtures occupied or traversed by animals.
3. Empty all troughs, racks, and other feeding and watering appliances.
4. Thoroughly clean all surfaces with soap or detergent, and rinse with water.
5. Saturate all surfaces with the recommended disinfecting solution for a period of 10 minutes.
6. Immerse all halters, ropes, and other types of equipment used in handling and restraining animals, as well as forks, shovels, and scrapers used for removing litter and manure.
7. Ventilate buildings, coops, cars, trucks, boats, and other closed spaces. Do not house animals or employ equipment until treatment has been absorbed, set, or dried.
8. After treatment with disinfectant, thoroughly scrub feed racks, troughs, automatic feeders, fountains, and waterers with soap or detergent, and rinse with potable water before reuse.

#### PRECAUTIONARY STATEMENTS

##### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**DANGER. CORROSIVE.** Causes irreversible eye damage and skin burns. Harmful if swallowed. Do not get in eyes, on skin, or on clothing. Wear protective eyewear (goggles, face shield or safety glasses), protective clothing and protective gloves (rubber or chemical resistant) Harmful if inhaled. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash clothing before reuse.

##### FIRST AID

**IF IN EYES:** Hold eye open and rinse slowly and gently with water for 15 - 20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.

**IF ON SKIN OR CLOTHING:** Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 - 20 minutes.

**IF SWALLOWED:** Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.

**IF INHALED:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible.

Call a poison control center or doctor for treatment advice.

Have the product container or label with you when calling a poison control center or doctor or going for treatment.

**NOTE TO PHYSICIAN:** Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsion may be needed.

##### STORAGE AND DISPOSAL

Store in original container in areas inaccessible to children. Do not reuse empty container. Wrap and discard in trash or recycle.

# KennelSol HC™

## Germicidal Detergent & Deodorant

**Disinfectant Bactericidal Fungicidal Mildewstatic \*Virucidal**

A multi-purpose, neutral pH, germicidal detergent and deodorant effective in hard water up to 400 ppm (calculated as CaCO<sub>3</sub>) in the presence of a moderate amount of soil (5% organic serum) according to the AOAC Use-dilution Test. Disinfects, cleans, and deodorizes in one labor-saving step.

Effective against the following pathogens:

(Bacteria:)	Serratia marcescens	*Respiratory Syncytial Virus (RSV)
Pseudomonas aeruginosa <sup>1</sup>	Shigella flexneri	(Animal Viruses:)
Staphylococcus aureus <sup>1</sup>	Shigella sonnei	*Avian polyomavirus
Salmonella choleraesuis	Staphylococcus aureus - Methicillin resistant (MRSA)	*Canine parvovirus - at 2 oz. per gallon
Acinetobacter calcoaceticus	Staphylococcus aureus - Vancomycin Intermediate Resistant (VISA)	*Canine distemper
Bordetella bronchiseptica	Staphylococcus epidermidis <sup>2</sup>	*Feline leukemia
Chlamydia psittaci	Streptococcus faecalis <sup>1</sup>	*Feline picornavirus
Enterobacter aerogenes	Streptococcus pyogenes	*Infectious bovine rhinotracheitis
Enterobacter cloacae	Streptococcus pyogenes	*Infectious bronchitis (Avian IBV)
Enterococcus faecalis - Vancomycin Resistant (VRE)	<sup>1</sup> ATCC & antibiotic-resistant strain	*Pseudorabies (PRV)
Escherichia coli <sup>1</sup>	<sup>2</sup> antibiotic-resistant strain only (Viruses:)	*Rabies
Fusobacterium necrophorum	*Adenovirus Type 4	*Transmissible gastroenteritis virus (TGE)
Klebsiella pneumoniae <sup>1</sup>	*Hepatitis B Virus (HBV)	(Fungi:)
Legionella pneumophila	*Herpes Simplex Type 1	Aspergillus niger
Listeria monocytogenes	*Herpes Simplex Type 2	Candida albicans
Pasteurella multocida	*Influenza A / Hong Kong	Trichophyton mentagrophytes
Proteus mirabilis	*Vaccinia	
Proteus vulgaris	*Rubella (German Measles)	
Salmonella enteritidis		
Salmonella typhi		
Salmonella typhimurium		



Manufactured for:

**Alpha Tech Pet, Inc.**

Lexington, Massachusetts 02420